

# Explaining the Gamma-Ray Burst $E_{peak}$ Distribution

Bradley E. Schaefer

*Department of Astronomy, University of Texas, Austin, TX 78712*

## ABSTRACT

The characteristic photon energy for Gamma Ray Bursts,  $E_{peak}$ , has a remarkably narrow distribution for bursts of similar peak flux, with values between 150 and 600 keV for most faint bursts. This result is surprising within the framework of internal shock models, since spectral shifts associated with the jet's blue shift (by a Lorentz factor of  $\Gamma$ ) and the cosmological red shift (by a factor of  $1+z$ ) should cause substantial smearing in the distribution of the spectral peak in the jet's co-moving frame,  $E_{rest}$ . For the general case where the luminosity ( $L$ ) varies as  $\Gamma^N$  and  $E_{rest}$  varies as  $\Gamma^M$ , then the observed  $E_{peak}$  will vary as  $L^{(M+1)/N}(1+z)^{-1}$ . For two independent set of 20 and 84 bursts,  $E_{peak}(1+z)$  varies as a power law of the luminosity with an index of  $(M+1)/N = 0.36 \pm 0.03$ . With this measured value, the above functional dependence of  $E_{peak}$  on  $L$  and  $z$  results in  $E_{peak}$  being roughly constant for bursts of similar peak flux,  $P_{256}$ . Thus, the kinematic smearing will be small, hence allowing the  $E_{peak}$  distribution to be narrow. This model also predicts that bright bursts will have high  $E_{peak}$  values because they all have some combination of high luminosity (and hence a large blue shift  $\Gamma$ ) and a nearby distance (and hence a small cosmological red shift). Quantitatively,  $E_{peak}$  should vary roughly as  $P_{256}^{0.36}$ , and this model prediction is strikingly confirmed with BATSE data by Mallozzi et al. A prediction of this model is that GRBs at very high red shift ( $z \sim 10$ ) should all appear with  $E_{peak}$  at  $\approx 200$  keV. A further prediction of this model is that normal bursts with  $P_{256}$  below the BATSE trigger threshold will appear as x-ray flashes with  $E_{peak} \approx 70$  keV; just as is reported by Kippen et al. and Heise et al.

*Subject headings:* gamma rays: bursts

## 1. Introduction

Gamma-Ray Bursts (GRBs) are indeed 'gamma-ray' bursts since their spectrum is dominated by gamma radiation.  $E_{peak}$  is the characteristic photon energy measured from a burst

spectrum by fitting a smoothly broken power law (Band et al. 1993). Mallozzi et al. (1995) have measured  $E_{peak}$  values for 399 BATSE bursts. They find that a histogram of  $E_{peak}$  is roughly log-normal in shape, with average values of  $\sim 200$  keV. They also find that the width of this distribution is surprisingly narrow, with a dispersion corresponding to a factor of  $\sim 2$  about the average. Brainard et al. (1998) have proven that the narrowness of this distribution is not an artifact of BATSE detector properties or of the analysis procedures, and in particular that the sharp falloff in frequency down to at least 50 keV is certain. Further, deep searches with the Solar Maximum Mission Gamma Ray Spectrometer have proven that the population of high- $E_{peak}$  bursts is nearly zero (Harris & Share 1998).

This is not to say that 'GeV bursts' and 'keV bursts' do not exist. For example, GRB950425 is certainly a GeV burst (Schaefer et al. 1998). Also, Kippen et al. (2001) and Heise et al. (2001) report x-ray transients that have similar light curves and durations as classical GRBs. The existence of these x-ray transients might be interpreted as implying that the  $E_{peak}$  distribution has a broad tail to low energies, despite the iron-clad data from BATSE. The resolution lies in the realization that the x-ray transients all have peak fluxes (in the 50-300 keV band) that are far below the normal BATSE trigger threshold. The result from Mallozzi et al. (1995) is that the  $E_{peak}$  distribution is narrow *for bursts in a small range of peak flux*. Bursts with peak fluxes far below the BATSE threshold do not (and should not) enter into the histograms constructed from BATSE triggers. If an  $E_{peak}$  distribution is constructed for very low peak flux events (including the x-ray transients), then a simple extrapolation of the average  $E_{peak}$  versus peak flux relation of Mallozzi suggests that such events will appear as ordinary bursts except with  $E_{peak} \sim 70$  keV, just as observed.

So apparently there is something about GRBs (the  $E_{peak}$  value) that is fairly constant over all bursts. This poses a double challenge to theory, since there must be some physical mechanism which acts as a thermostat and there must be some mechanism for balancing the kinematic blue shifts (due to the jet's velocity towards Earth) and red shifts (due to the cosmological expansion). Based on the factor of  $\sim 1000$  variation in burst luminosity, the Lorentz factor of the jet likely varies by a large factor, while the cosmological red shift varies by over a factor of 3; so we might expect the kinematic effects to produce a broad  $E_{peak}$  distribution even if all bursts had the same rest frame temperature.

Mallozzi et al. (1995) also found that the average  $E_{peak}$  value varied systematically with the apparent brightness of the burst. The peak flux used here,  $P_{256}$ , is measured over a 256 ms time scale within a limited energy band from 50-300 keV (Fishman et al. 1994). The bright BATSE bursts have  $\langle E_{peak} \rangle$  around 339 keV while the faintest BATSE bursts have  $\langle E_{peak} \rangle$  around 175 keV. The trivial explanation of this effect as arising only from cosmological red shifts is unreasonable since burst peak fluxes vary more due to the wide

spread in luminosity than due to distances. So a second mystery is why  $\langle E_{peak} \rangle$  varies with  $P_{256}$  as observed.

## 2. The Narrow $E_{peak}$ Distribution

Two kinematic effects will change the characteristic photon energy of the emitting region in its own rest frame,  $E_{rest}$ , into the characteristic photon energy as seen by Earth,  $E_{peak}$ . The first is the blue shift caused by the bulk velocity of the emitting region towards Earth. This will increase the detected characteristic energy by a factor of  $\Gamma$ . The second is the red shift caused by the usual cosmological expansion of the Universe. This will decrease the detected characteristic energy by a factor of  $(1+z)^{-1}$ , where  $z$  is the red shift. So,

$$E_{peak} = E_{rest} \Gamma (1+z)^{-1}. \quad (1)$$

Here, the  $E_{peak}$  distribution will be a convolution of the  $E_{rest}$  distribution and the kinematic smearing effects from the  $\Gamma(1+z)^{-1}$  factor.

The  $E_{rest}$  value can vary from pulse-to-pulse and burst-to-burst. In general,  $E_{rest}$  might have a power law dependence on  $\Gamma$ . So

$$E_{rest} = E_0 \Gamma^M, \quad (2)$$

where  $M$  is the power law index. The exponent  $M$  might be zero if there is some process which limits the temperature in the emitting region, or  $M$  might be unity if the kinetic energy of the jet is entirely converted to thermal energy and then into gamma radiation.

Within generic internal shock scenarios, the burst isotropic luminosity,  $L$ , varies as some power of  $\Gamma$ . One power of  $\Gamma$  comes from the available energy in the shell. Two more powers of  $\Gamma$  come from the blue shifting of photons in the usual  $E^{-2}$  burst spectrum. (The 50-300 keV power law index varies over a typical range of about unity in width from burst-to-burst, with an average of close to -2 as shown in Fig. 4 of Schaefer et al. (1998) and Fig. 46 of Schaefer et al. (1994).) The bunching of photons by the forward motion of the shell will not increase the luminosity since the overall timing of sub-pulses (and thus the distribution of photons within the peak) is determined by the central engine which has no  $\Gamma$  dependence. (Alternatively, if a single sub-pulse with observed duration longer than 256 ms dominates the peak flux, then an additional  $\Gamma^2$  dependence is due to the bunching of photons.) Hence, the luminosity should vary as something like  $\Gamma^3$ . However, variations on this basic scenario are possible; for example the peak luminosity might well be dominated by a relatively small emission area (to account for the fact that most bursts have millisecond variations (Walker, Schaefer, & Fenimore 2000)) and the resulting relativistic beaming could provide another

factor of  $\Gamma^2$ . Emission efficiencies could well have some dependency on  $\Gamma$ . To allow for any reasonable scenario, let me adopt a general expression of

$$L = L_0 \Gamma^N, \quad (3)$$

where  $N$  is the power law index that is expected to be  $\sim 3$  or perhaps  $\sim 5$  and to vary somewhat from burst-to-burst.

Equations 1-3 can be combined as

$$E_{peak} = E_0 (L/L_0)^{(M+1)/N} (1+z)^{-1}. \quad (4)$$

The kinematic part can be isolated as

$$E_{peak} \propto L^{(M+1)/N} (1+z)^{-1}. \quad (5)$$

This representation is convenient since  $L$  and  $z$  can be measured for many bursts, either by direct measures of their optical red shifts or based on the lag and variability distance indicators (Schaefer, Deng, & Band 2001). This is also convenient since lines of constant  $P_{256}$  are also functions of  $L$  and  $z$ , and the narrow  $E_{peak}$  distributions of Mallozzi et al. (1995) were constructed for bursts in narrow bins of  $P_{256}$ . Figure 1 displays curves of constant  $E_{peak}$  (for the case of  $(M+1)/N = 0.36$ ) and lines of constant  $P_{256}$ . The two  $P_{256}$  lines are for values  $(10.0 \text{ and } 1.0 \text{ ph s}^{-1} \text{ cm}^{-2})$  which are the medians of Mallozzi's brightest and dimmest bins (for  $\Omega_M = 0.3, \Omega_\Lambda = 0.7, H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ). The two curves of constant  $E_{peak}$  values (339 and 175 keV) were chosen to correspond to the measured values for Mallozzi's brightest and dimmest bins (for  $L_0$  and  $E_0$  normalized so that the lower curve is close to the faintest  $P_{256}$  line).

Over the range for which GRBs are seen (typically  $0.4 < z < 4$ ), the  $E_{peak}$  curves are fairly close to the  $P_{256}$  lines. This is to say that for a collection of bursts within a narrow  $P_{256}$  bin, the kinematic shifts do not vary much. Over a wide range of  $L$  or  $z$ , the overall shifting of  $E_{rest}$  is roughly constant. This constancy is an essential compensation between the red and blue shifts, in that bursts with low blue shifts (and hence low luminosities) must necessarily be close to Earth and have offsetting low red shifts; while the high blue shift jets (with high  $L$ ) are distant with an offsetting high red shift. So the combined kinematic smearing varies little, thus allowing GRBs to have a narrow  $E_{peak}$  distribution despite having a wide range of red shifts and luminosities.

A narrow  $E_{peak}$  distribution requires both that the kinematic smearing be small and that the  $E_{rest}$  distribution be at least as narrow. The explanation just given answers only part of the requirements. A physical mechanism to explain why the  $E_{rest}$  distribution is so narrow is still needed (such as in Kazanas, Georganopoulos, & Mastichiadis 2002).

The size of the kinematic smearing depends on the exponent of  $L$  in Eq. 5. There will be additional smearing due to the finite width of the  $P_{256}$  bins. With the model from Eq. 5, I have calculated the total smearing across the bin for the faintest bursts ( $0.95 < P_{256} < 1.3$ ) over the normal range of red shifts ( $0.4 < z < 4$ ). For an acceptable value of  $(M+1)/N$ , the smearing across this bin should be smaller than the observed spread of  $E_{peak}$ . The FWHM of the  $E_{peak}$  distribution for the faintest bursts stretches from roughly 150 keV to 600 keV, a factor of 4 (Mallozzi et al. 1995). The kinematic smearing is less than a factor of 4 for all values of  $(M+1)/N$  less than 0.45. The least smearing (at 19%) occurs for  $(M+1)/N = 0.21$ .

### 3. How $E_{peak}$ Varies With $P_{256}$

Nemiroff et al. (1994) discovered that bright bursts are harder than faint bursts, while Mallozzi et al. (1995) showed specifically how  $\langle E_{peak} \rangle$  varies with  $P_{256}$  (see Fig. 2). The general formalism from the previous section can yield a prediction for how  $\langle E_{peak} \rangle$  varies with  $P_{256}$ . The reason for this variation is a consequence of Eq. 5. From Fig. 1, we see that there is nearly a one-to-one correspondence between  $\langle E_{peak} \rangle$  and  $P_{256}$ . But the exact predicted function will depend on the distribution of bursts within a  $P_{256}$  bin. If the median burst is roughly at a constant red shift (specifically, at the strong inflection in the burst number density at  $z \approx 1$  (Schaefer, Deng, & Band 2001)), then Eq. 5 becomes

$$E_{peak} \propto P_{256}^{(M+1)/N}, \quad (6)$$

since  $L \propto P_{256}$ . Alternatively, if the median burst is roughly at a constant luminosity (specifically, at the break in the GRB luminosity function around  $2 \times 10^{52} \text{ erg s}^{-1} \text{ cm}^{-2}$  (Schaefer, Deng, & Band 2001)), then Eq. 5 takes on a substantially more complex functional form that is similar to Eq. 6. The likely case is midway between these two extreme alternatives, with the median burst in each  $P_{256}$  bin along a line roughly perpendicular to the line of constant  $P_{256}$ . Here, there is no simple expression although the  $E_{peak}$  distribution is close to a power law in  $P_{256}$ .

The  $\langle E_{peak} \rangle$  data from Mallozzi et al. (1995) is displayed in Figure 2. What value of  $(M+1)/N$  best fits the data based on Eq. 6? The chi-square is minimized for  $(M+1)/N$  equal to  $0.40 \pm 0.14$ . Here, the median burst in each  $P_{256}$  bin is taken to lie on a line perpendicular to the line of constant  $P_{256}$  in the  $L$  versus  $z$  plot. In the next section, I will show that  $(M+1)/N$  is equal to  $0.36 \pm 0.03$ , and this is still an acceptable fit (see Fig. 2).

The point of this section is that Eq. 5 provides a simple and general explanation for the observed variation of  $\langle E_{peak} \rangle$  with  $P_{256}$  (Eq. 6). In words, the brighter bursts will have some combination of higher luminosities (with a higher  $\Gamma$ ) and be closer (with a smaller  $1+z$ )

than fainter bursts, so that the jet’s blue shift and cosmological red shift will work together to give a higher  $E_{peak}$  for the brighter bursts.

#### 4. The Values of $M$ and $N$

The values of  $M$  and  $N$  can tell us about the physics of gamma ray bursts. Observations can place reliable constraints on both  $M$  and  $N$ . Sections 2 and 3 have already provided limits, and two more observational constraints will now be presented.

For the 112 BATSE bursts with known  $L$  and  $z$  (based on the measured lag and variability) reported in Schaefer, Deng, & Band (2001), the  $E_{peak}$  value at the time of peak flux is known for 84 events. (The reason for why the 28 events do not have a measured  $E_{peak}$  is not known due to the untimely death of R. Mallozzi.) Eq. 5 shows that  $E_{peak}(1+z)$  should vary with  $L$  as a power law with index  $(M+1)/N$ . As shown in Figure 3, the logarithms of  $E_{peak}(1+z)$  and  $L$  are indeed accurately related linearly. A chi-square fit gives  $(M+1)/N = 0.36 \pm 0.03$ .

This same method can be applied to 20 bursts which have the luminosities based on optically measured red shifts. Figure 3 shows these bursts, with the data taken from the literature (primarily Schaefer (2002) and Amati et al. (2002)). This sample also shows a power law dependence, with a highly significant correlation ( $r = 0.90$ , for a chance probability of  $3 \times 10^{-8}$ ). The slope is  $(M+1)/N$ , which is fitted to be  $0.38 \pm 0.11$ .

The variability/luminosity relation (Fenimore & Ramirez-Ruiz 2000) can be used to constrain  $N$ . Variability ( $V$ ) is a measure of how ‘spikey’ a burst light curve is, which empirically varies as the inverse of the rise times in the light curves. The rise times will be limited by the geometric delays which scale as  $\Gamma^{-2}$ , so  $V \propto \Gamma^2$ . Schaefer (2002) has shown from 9 bursts with measured red shifts that  $L \propto V^{1.57 \pm 0.17}$ . Thus, we have  $L \propto \Gamma^{3.14 \pm 0.34}$ , which is to say  $N = 3.14 \pm 0.34$ .

I now have four observational restrictions and two theoretical restrictions on  $N$  and  $M$ . (1) The narrowness of the observed  $E_{peak}$  distribution requires  $(M+1)/N < 0.45$ . (2) The variation of  $\langle E_{peak} \rangle$  with  $P_{256}$  is easily explained if  $(M+1)/N = 0.40 \pm 0.14$ . (3) The variation of  $E_{peak}(1+z)$  with  $L$  for two independent set of bursts shows that  $(M+1)/N$  equals  $0.36 \pm 0.03$  and  $0.38 \pm 0.12$ . (4) The variability/luminosity relation gives  $N = 3.14 \pm 0.34$ . (5) Ordinary relativistic effects for any internal shock model implies that  $N \gtrsim 3$ . (6) The conversion of the jet energy to thermal energy suggests that  $0 \geq M \geq 1$ . Constraints 1-3 reduce to  $(M+1)/N = 0.36 \pm 0.03$ . Constraints 4 and 5 give  $N = 3.14 \pm 0.34$ . These two results then imply that  $M = 0.13 \pm 0.15$ , which is consistent with constraint 6. In round

numbers,  $N = 3$  and  $M = 0$ .

## 5. Implications

What should very high red shift ( $z \sim 10$ ) GRBs look like? A first guess (e.g., Bloom et al. 2001) might be that the cosmological red shift will transform them to appear more like x-ray bursts with  $E_{peak} \sim 20$  keV. The analysis in this paper shows that this first guess is wrong since the  $z \sim 10$  bursts must have a very high luminosity and hence a very high  $\Gamma$  which will blue shift the  $E_{peak}$  back to gamma-ray energies. In particular, from either equation 5 or Figure 1, we see that the highest red shift bursts will all have the same  $E_{peak}$  as the nearby events of the same  $P_{256}$ . Thus, any  $z > 5$  bursts that are in the BATSE catalog will have  $E_{peak} \sim 200$  keV.

Kippen et al. (2001) have used BATSE real-time data to look at fast x-ray transients discovered by the BeppoSAX satellite (Heise et al. 2001). They find that these events have similar light curves and durations as GRBs, which is suggestive that the events are normal bursts. Their peak flux is up to 3 times lower than the BATSE trigger threshold while their median  $E_{peak}$  value is 70 keV. The natural suggestion was made that very faint and very soft bursts might be at very high red shift. However, the model presented in this paper shows that *all* bursts below the BATSE threshold will on average have a low  $E_{peak}$ , with no necessity of high red shift. (This is also realized from a simple extrapolation of the data from Mallozzi et al. as shown in Figure 2.) That is, a low  $P_{256}$  is due to some combination of low luminosity (hence a low blue shift from the jet) and large distance (hence a large cosmological red shift) which will produce systematically low  $E_{peak}$  values. While some of the fast x-ray transients might be at high  $z$ , the steepness of the GRB luminosity function implies that almost all are at moderate red shift.

It is disappointing that lines of constant  $E_{peak}$  are closely parallel to lines of constant  $P_{256}$  in the  $L$  versus  $z$  plot. If this had not been true, then a simple measurement of  $E_{peak}$  and  $P_{256}$  would define the bursts' position in the plot and we would hence know the burst luminosity and red shift.

The value of  $E_0$  is approximately constant for all bursts. This constancy is similar to recent results that bursts are 'standard candles' from the lag/luminosity and variability/luminosity relations as well as that the total energy of bursts is nearly a constant. Thus it now seems that many aspects of GRBs are constant, despite the chaotic appearance of their light curves. One implication of this constancy is that GRBs might prove useful as tools for cosmology.

The lag/luminosity and variability/luminosity relations will be the primary methods of getting luminosity distances. Unfortunately, these relations are not as tight as we might hope. It is possible that  $E_{peak}$  might be used as a third parameter to improve the distance indicators, much as the Cepheid luminosities are improved with a period/color/luminosity relation. Thus, lag/ $E_{peak}$ /luminosity and variability/ $E_{peak}$ /luminosity relations might substantially improve the accuracy of GRBs as standard candles.

I thank Robert Mallozzi and the BATSE Team for providing the  $E_{peak}$  values for the 84 bursts.

## REFERENCES

- Amati, L. et al. 2002, A&A, 390, 81
- Band, D. L. et al. 1993, ApJ, 428, 21
- Bloom, J. S., Frail, D. A., & Sari, R. 2001, AJ, 121, 2879
- Brainard, J. J. et al. 1998, 20th Texas Symposium, Paris (astro-ph/09904039)
- Fenimore, E. E., & Ramirez-Ruiz, E. 2000, ApJ, submitted, (astro-ph/0004176)
- Fishman, G. J. et al. 1994, ApJS, 92, 229
- Harris, M. J. & Share, G. H. 1998, ApJ, 494, 724
- Heise, J., in't Zand, J., Kippen, R. M., & Woods, P. M. 2001, in Gamma-Ray Bursts in the Afterglow Era, eds E. Costa, F. Frontera, and J. Horth (Berlin; Springer), p. 16 (astro-ph/0111246)
- Kazanas, D., Georganopoulos, M., & Mastichiadis, A. 2002, ApJ, 578, L15
- Kippen, R. M., Woods, P. M., Heise, J., in't Zand, J., Preece, R. D., & Briggs, M. S. 2001, in Gamma-Ray Bursts in the Afterglow Era, eds E. Costa, F. Frontera, and J. Horth (Berlin; Springer), p. 22 (astro-ph/0102277)
- Mallozzi, R. S., et al. 1995, ApJ, 454, 597
- Nemiroff, R. J. et al. 1994, ApJ, 435, L133
- Schaefer, B. E. 2002, ApJ, submitted



Schaefer, B. E., Deng, M., & Band, D. L. 2001, *ApJ*, 563, L123

Schaefer, B. E., J. Hobbeheydar, Bromm, V., & Fenimore, E. 1998, in *Gamma Ray Bursts*, eds. C. A. Meegan, R. Preece, and T. Koshut (New York, AIP), p. 379

Schaefer, B. E. et al. 1994, *ApJSupp*, 92, 285

Schaefer, B. E. et al. 1998, *ApJ*, 492, 696

Walker, K. C., Schaefer, B. E., & Fenimore, E. E. 2000, *ApJ*, 537, 264




Fig. 1.— Lines of constant  $E_{peak}$  and  $P_{256}$ . In a plot of luminosity versus red shift, bursts of constant peak flux will fall on a nearly straight line. Plotted are two dashed lines for  $P_{256}$  equal to 1.0 and 10.0  $photons\ cm^{-2}\ s^{-1}$ , which corresponds to the center of the faintest and brightest bins used by Mallozzi et al. (1995). Also plotted are two curves of constant  $E_{peak}$ , as taken from Eq. 5 for  $(M + 1)/N = 0.36$ . The normalization factor has been chosen such that the lower curve with  $E_{peak} = 175\ keV$  (corresponding to the faintest bursts) overlaps the  $P_{256}$  bin for the faintest bursts. The upper curve is calculated for  $E_{peak} = 339\ keV$ , which corresponds to Mallozzi’s brightest bursts. The main point from this figure is that the lines and curves are roughly parallel over the range of observed bursts (roughly  $0.4 < z < 4$ ), and this means that the blue shifts (from the jet’s motion towards Earth) largely cancels out the red shifts (from the Universal expansion) for bursts of similar peak flux. The curves and lines are not perfectly parallel which implies that kinematic effects will contribute to some of the observed scatter in the  $E_{peak}$  distribution.




Fig. 2.— The model  $E_{peak}$  as a function of  $P_{256}$ . Mallozzi et al. (1995) showed how the average  $E_{peak}$  changes systematically with  $P_{256}$ , as shown by the plotted data in the figure. From Fig. 1, we see that the model predicts a fairly close correspondence between  $E_{peak}$  and  $P_{256}$ , with  $E_{peak}$  increasing with  $P_{256}$ . From Eq. 6, for the  $(M + 1)/N = 0.36$  case with an arbitrary normalization, the model prediction is shown as a smooth curve. The reason for the variation is that bright bursts are on average of higher luminosity (with larger  $\Gamma$  increasing the blue shift of the jet) and of nearer distance (with smaller  $z$  decreasing the cosmological red shift) than for fainter bursts. There is a good match between model and observations.




Fig. 3.— Direct fit of Eq. 5 to  $E_{peak}$  data. The quantity  $E_{peak}(1+z)$  is related as a power law to the burst luminosity, and the slope will give the value for  $(M+1)/N$ . This is shown here for two independent data sets for which the luminosities are derived by two independent methods. The first data set is for 20 bursts with spectroscopically measured red shifts (the open circles). The second is for 84 bursts (whose binned values are shown as filled diamonds) whose luminosity (and then red shift) were determined with the spectral lag and variability light curve parameters. Both data sets show a highly significant and similar power law relation. The existence of this relation demonstrates the validity of equation 5. The explanation for this empirical relation is that the high luminosity bursts have a high  $\Gamma$  for their jet (which is why they are of high luminosity) which then blue shifts the  $E_{rest}$  value to a high observed  $E_{peak}$  value. The fitted values for  $(M+1)/N$  are  $0.36 \pm 0.03$  and  $0.38 \pm 0.11$  respectively.